

第一著者の観察ならびに採集者の一人、城戸克弥氏からの情報をもとに、解剖を行った結果、本種は、軟体のヤスデ類イソフサヤスデを食べているという特異な性質が、明らかになった。

References

- HABU, A., 1967. Carabidae: Truncatipennes group (Insecta: Coleoptera). *Fauna Japonica*. ii + xiv + 338 pp., xxvii pls. Tokyo Electrical Engineering College Press, Tokyo.
- 1982. Revised and supplementary notes on and descriptions of the Truncatipennes group of Japan (I) (Coleoptera, Carabidae). *Ent. Rev. Japan, Osaka*, **6**: 85–142.
- 1983. Ditto (III). *Ibid.*, **38**: 105–146.
- HOPE, F. W., 1838. The coleopterist's manual, part the second, containing the predaceous land and water beetles of LINNEUS and FABRICIUS. xvi + 168 pp., 3 pl. London, Bohn. frontispiece.
- JEDLIČKA, A., 1963. Monographie des Truncatipennen aus Ostasien Lebiinae – Odacanthinae – Brachyninae (Coleoptera, Carabidae). *Ent. Abh. Mus. Tierk. Dresden*, **32**: 269–579, 6 col. pls.

Elytra, Tokyo, **38**(1): 93–94, May 31, 2010

Records of Some Nanophyid Weevils (Coleoptera, Nanophyidae) New to Laos, Part 2

Junnosuke KANTOH and Hiroaki KOJIMA

Laboratory of Entomology, Tokyo University of Agriculture,
1737 Funako, Atsugi, Kanagawa, 243-0034 Japan

The nanophyid weevil fauna of Laos has poorly been studied and merely a total of two genera and three species are known from the country (KANTOH & KOJIMA, 2009 b). In this paper, two genera and four species are newly added to the fauna based on the recent survey conducted at rice paddy and its adjacent areas.

We thank Mr. H. WAKAHARA and his family for their arrangement of our trip. Thanks are due to Dr. S. OKAJIMA for his encouragement, and Ms. M. WATANABE, Messrs. T. BAN and G. OISHI for their help in the field. This study is supported by the KAKENHI (21405019, head investigator: S. OKAJIMA).

1. *Alonsiellus pubescens* (ROELOFS, 1874)

Specimen examined. 1 female, Ban Don (alt. ca. 1,100 m, rice paddy), Houapan Prov., 25–VIII–2009, G. OISHI leg.

Distribution. Japan, E. Siberia, Korea, Laos. New to Laos.

Weevils are associated with *Ludwigia epilobioides* (Onagraceae) inhabiting humid places and

make the stem gall in Japan (Y. SAWADA, pers. comm.).

2. *Nanophyes albovittatus* ROELOFS, 1874

Specimens examined. 1 male, Ban Don (alt. ca. 1,100 m, rice paddy), Houapan Prov., 16–VI–2009, T. BAN leg. 3 males and 1 female, Muang Kham Hot Spring (alt. 626 m), N 19° 33' / E 103° 26', Xieng Khouang Prov., 20–VIII–2009, H. KOJIMA leg.

Distribution. Japan, Korea, Laos. New to Laos.

Weevils are associated with *Ludwigia epilobioides* and make the fruit gall in Japan.

3. *Nanophyes pallipes* ROELOFS, 1874

Specimens examined. Houapan Prov.: 2 males and 5 females, Ban Nasala (alt. ca. 1,300 m, rice paddy), 15–VI–2009, H. KOJIMA leg. 1 female, Ban Don, (alt. ca. 1,100 m, rice paddy), 25–VIII–2009, H. KOJIMA leg. Khammouan Prov.: 1 male, 7 km east from Thakhek (alt. 169 m), 18–II–2010, H. KOJIMA leg. 2 males and 1 female, 13 km east from Thakhek (alt. 188 m), 18–II–2010, H. KOJIMA leg.

Distribution. Japan (incl. Ryukyus), Laos. New to Laos.

Weevils are associated with *Rotala indica* (Lythraceae) and make the stem gall.

4. *Shiva trispinosus* PAJANI et BHATEJA, 1982

Specimens examined. Khammouan Prov.: 6 males and 6 females, Thakhek (alt. 185 m), 19–IX–2009, G. OISHI leg. (on flower of *Lagerstroemia* sp.). 2 males and 1 female, 7 km east from Thakhek (alt. 169 m), 18–II–2010, T. BAN, H. KOJIMA, M. WATANABE & J. KANTOH leg. Vientiane Dist.: 1 female, Tha Ngon (alt. 254 m), 29–IX–2009, H. WAKAHARA leg. Salavanh Prov.: 1 male, 94 km north from Pakse (alt. 182 m), 12–II–2010, H. KOJIMA leg.

Distribution. India, Indonesia (Bali), Laos. New to Laos.

Weevils are associated with crape myrtle, *Lagerstroemia* spp. (Lythraceae) (KANTOH & KOJIMA, 2009 a).

References

- KANTOH, J., & H. KOJIMA, 2009 a. Records of the nanophyid weevils (Coleoptera, Nanophyidae) from Bali, Indonesia. *Elytra, Tokyo*, **37**: 171–174.
 ——— 2009 b. Records of some nanophyid weevils (Coleoptera, Nanophyidae) new to Laos. *Ibid.*, **37**: 373–374.